# COMMENT

# Where is WAIR (and other wing-assisted behaviours)? Essentially everywhere: a response to Kuznetsov and Panyutina (2022)

ASHLEY M. HEERS<sup>1</sup>, BRET W. TOBALSKE<sup>2</sup>, BRANDON E. JACKSON<sup>3</sup> and KENNETH P. DIAL<sup>2,\*,•</sup>

<sup>1</sup>Department of Biological Sciences, California State University Los Angeles, Los Angeles, CA 90032, USA <sup>2</sup>Flight Laboratory, Field Research Station at Fort Missoula, Division of Biological Sciences, University of Montana, Missoula, MT 59804, USA <sup>3</sup>Department of Biological and Environmental Sciences, Longwood University, Farmville, VA 23909, USA

Received 31 March 2022; revised 27 May 2022; accepted for publication 30 May 2022

Kuznetsov and Panyutina (2022) offer a reanalysis of the kinematic and force plate data previously published by Bundle and Dial (2003). Their intention is to describe instantaneous wing forces during wing-assisted incline running (WAIR), focusing particularly on the upstroke phase. Based on their interpretation of wing forces and muscle function, the authors conclude that 'WAIR is a very specialized mode of locomotion that is employed by a few specialized birds as an adaptation to a very specific environment and involving highly developed flying features of the locomotor apparatus', and thus not relevant to the evolution of avian flight. Herein, we respond to the authors' interpretations, offering an alternative perspective on WAIR and, more generally, on studies exploring the evolution of avian flight.

ADDITIONAL KEYWORDS: cooperative use of wings and legs - evolution of avian flight.

#### INTRODUCTION

We thank Kuznetsov and Panyutina (2022; hereafter, the 'authors'), and many other researchers, for their interest in wing-assisted incline running (WAIR). Here we reiterate points raised in our submitted review and respond to the main conclusions presented by the authors. A novel contribution of their analysis is to describe the instantaneous wing forces during the upstroke, and we do not question their key conclusion that forces are produced by the wings during the upstroke. However, as we describe, these forces are more likely to be inertial than aerodynamic. We also propose that musculoskeletal activity during WAIR is actually reasonably consistent with the evolutionary inferences summarized by the authors.

Based on their analyses, the authors suggest that the upstroke of an adult chukar engaged in WAIR is a unique, specialized aerodynamic event among birds, and that it should be regarded as a crown locomotor specialization not suitable as a model for the evolutionary origins of flight. The authors' rejection of WAIR as an analogue to the early evolution of avian wings is based on aerodynamic and musculoskeletal considerations, which we address, with contrary conclusions. For the purposes of discussion, we will focus on 'variant B' of the authors' reanalysis of data in Bundle & Dial (2003).

# AERODYNAMIC CONSIDERATIONS

Based on a recalculation of data from Bundle & Dial (2003), the authors state that the average force acting on the wings is 113% body weight (variant B), that in the sagittal plane this force is primarily an aerodynamic force, and that the force is directed into the substrate and upward during the middle and end of the downstroke and downward during the rest of the wingbeat cycle. Averaged over the entire wingbeat

© The Author(s) 2022. Published by Oxford University Press on behalf of The Linnean Society of London. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com

<sup>\*</sup>Corresponding author. E-mail: kpdial@gmail.com

cycle, the force is directed 11° below perpendicular to the substrate surface for variant B. That the force is directed partially toward the substrate is consistent with previous work on WAIR and not a new discovery; indeed, Dial (2003) describes the wings as 'acting like the spoilers on a race car to improve traction' during WAIR (Bundle & Dial, 2003; Dial et al., 2006, 2008; Tobalske & Dial, 2007). However, if the analysis is correct, and if the large force generated during the upstroke is an aerodynamic force as the authors suggest, then this finding contrasts with every previously published study on upstroke aerodynamics in bird flight, as outlined below.

To begin, it is first important to note that **the tip**reversal upstroke used by an adult chukar is not unique or particularly specialized among birds. Light and X-ray video (Baier et al., 2013; Heers et al., 2016) show that chukars use relatively similar kinematics to achieve different behaviours, with joint movements becoming more exaggerated as the difficulty of the behaviour increases and culminating in an exaggerated tip-reversal upstroke. This upstroke style is widespread among avian species during movement at low advance ratios (J), where wholebody translational velocity is significantly less than wingtip velocity. For example, tip-reversal upstrokes are routinely used at low J in many birds that have relatively pointed, high-aspect ratio wings, such as parakeets and cockatiels (Tobalske et al., 2003), pigeons and doves (Tobalske, 2000), and dabbling ducks (Vazquez, 1994). Species with rounded wings typically use a flexed-wing upstroke (Tobalske, 2000), but phasianids, including the chukar, use a tip-reversal upstroke in spite of having relatively rounded wings (Tobalske & Dial, 2000). The kinematics of these tipreversal upstrokes have been recognized since (Marey, 1890). Additional kinematics are available in Lorenz (1933), Brown (1953), Tobalske & Dial (2000), Crandell & Tobalske (2015), and skeletal adaptations that facilitate hand-wing supination during tip-reversal upstrokes are in Vazquez (1992). In short, tip-reversal upstrokes are well known, widespread and fairly well studied.

Second, there is no evidence that any bird species operating at low advance ratios can produce aerodynamic force during the upstroke that is on par with force produced during the downstroke. Since Marey (1890), researchers have hypothesized that tip-reversal upstrokes provide substantial lift, but all available evidence instead suggests that aerodynamic force production is much less than in downstrokes. This should not be surprising given that hummingbirds, the only species to supinate almost their entire wing during the upstroke, have upstrokes that produce ~30% of downstroke lift (Warrick *et al.*, 2005, 2009). Several lines of evidence suggest that tip-reversal and flexed-wing upstrokes also contribute significantly less lift compared with downstrokes. For example, analysis of 3D kinematics and body mass distribution suggests that aerodynamic forces during tip-reversal upstrokes in turning pigeons are ~50% of those produced during downstrokes (Ros et al., 2011). Dried wings mounted in a mid-upstroke tip-reversal posture and spun like a propeller produce lift that is ~36% of the body weight for both wings (Crandell & Tobalske, 2011), although this likely overestimates in *vivo* function because the wings morph dynamically during upstroke, and full supination of the hand wing occurs only during the middle of upstroke (Crandell & Tobalske, 2011). Measures of the near wake using particle image velocimetry (PIV) in diamond doves (Crandell & Tobalske, 2011) demonstrate that the primary aerodynamically active portion of tipreversal upstroke occurs at the upstroke-downstroke transition, when the wings clap and peel. This has two effects: generating a thrusting impulse that is  $\sim 11\%$ of the downstroke impulse, and initiating an earlier onset of circulation (hence lift) on the wing during the downstroke. Accelerometry in cockatiels shows that the tip-reversal upstroke produces ~14% of the force of the downstroke (Hedrick et al., 2004), and measures of parrotlets in an aerodynamic force chamber reveal peak upstroke forces that are 10-15% of the downstroke, except during the final wingbeat of landing when the bird is using drag to decelerate (Chin & Lentink, 2017). Flexed-wing upstrokes likely produce even less aerodynamic force: zebra finches using this type of upstroke transmit minimal momentum to the upstroke wake compared to tip-reversal upstrokes (Crandell & Tobalske, 2015). Taken together, these studies suggest that the aerodynamic contribution of the downstroke far exceeds that of the upstroke.

The analysis of the authors motivated us to revisit our previous samples of PIV from chukars engaged in WAIR (Tobalske & Dial. 2007). Consistent with the data from the upstroke in diamond doves (Crandell & Tobalske, 2011), we observed induced air velocities from the late upstroke, immediately prior to the upstroke-downstroke transition. These velocities were less than those observed during the downstroke, and directed toward the substrate and not downward. In the example shown in Figure 1, the average induced velocity in the middle of the downstroke wake was 9.0 m s<sup>-1</sup>, oriented at 54° relative to horizontal, meaning that it was directing force upward and toward the ramp, which was angled at 90° (i.e. vertical). The area of the wake from the beginning of the upstroke overlapped with the downstroke-upstroke transition, producing a complex flow field (Tobalske & Dial, 2007). This early phase of upstroke did direct force down and away from the substrate (with an orientation of -54° relative to horizontal); however, the average velocity



**Figure 1.** Induced velocities in the wake of an adult chukar using wing-assisted incline running (WAIR) to ascend a ramp set at an incline of 90°. See Tobalske & Dial (2007) for methods. The approximate boundaries are indicated as closed loops. The boundaries overlap at the downstrokeupstroke transition.

during this phase was only  $0.8 \text{ m s}^{-1}$ . Late upstroke induced velocity was  $5.0 \text{ m s}^{-1}$  oriented at  $62^{\circ}$  relative to horizontal—slightly more upward than the orientation of the downstroke. A different example from an adult chukar flap-running up an  $80^{\circ}$  incline exhibited a late upstroke induced velocity of  $5.4 \text{ m s}^{-1}$  oriented at  $20^{\circ}$ relative to horizontal, thus more directed as thrust toward the substrate and less as weight support. These patterns are consistent with previous studies showing that upstroke aerodynamic force production is less than that of downstroke, and they suggest that such forces direct the bird toward the substrate and primarily upward (rather than downward).

Previously, we measured circulation and modelled projected wake area, and spun dried wings like propellers, to estimate the average force produced during the downstroke in adult chukars (Tobalske & Dial, 2007; Heers et al., 2011). These measurements indicated that aerodynamic force was approximately 60% of body weight during 65° WAIR. At a 70° incline, as in Bundle & Dial (2003), this force was likely somewhat higher. Nevertheless, even if the upstroke produced as much force as the downstroke-which seems unlikely for the reasons described above-peak aerodynamic forces are only generated for part of the wingstroke cycle, such that the aerodynamic force produced over the entire wingbeat would probably be far less than 60% body weight. This is substantially lower than the 113% calculated by the authors.

Based on these points, we instead suggest that inertia is contributing substantially to the force acting on the wings. The authors state that inertial forces 'should be mainly confined to the wingbeat plane' and that they 'cannot influence significantly the 2D accelerometer system in the sagittal-bound

experimental setting'. This conclusion assumes (1) that peak magnitudes of circumferential inertial force occur at the downstroke-upstroke and upstroke-downstroke transitions, when the wing approaches the sagittal plane and circumferential inertial force would be directed transversely, and (2) that inertial forces during the rest of the wingbeat cycle are very small. However, simulations suggest that inertial forces are quite substantial through much of the wingbeat cycle (Heers et al., 2018), and negligible only during the mid-downstroke and the mid-upstroke, indicating that inertia would contribute to forces acting on the wings during early and late downstroke and upstroke. Any upstroke with a partially extended wing requires inertial work from the wing muscles to move the wing. Indeed, direct measures of contractile behaviour of the supracoracoideus muscle in pigeons using tip-reversal upstrokes during slow flight reveal that the inertial power required for the upstroke is within one standard deviation of the power output by the muscle (Tobalske & Biewener, 2008).

In addition, the wingstroke plane is based on the path of the wingtip through space. However, the mass of the wing lies posterior to the wingtip and leading edge of the wing, because the humerus is retracted [particularly during WAIR (Heers et al., 2016, 2018)] and the elbow is bent, such that the mass of the wing muscles, largest wing bones (humerus, radius, ulna) and feathers (whose mass influences inertial calculations) lies posterior to the leading edge and stroke plane. These masses thus rotate about the shoulder joint and likely exert torques that do not operate perpendicular to the sagittal plane of the bird, which should, in turn, result in inertial forces that are partially directed either anteriorly or posteriorly, depending on the portion of the wingbeat cycle. Indeed, simulations of WAIR without any aerodynamic force production result in substantial joint moments through most of the wingbeat cycle, including moments along the vertebral axis, suggesting that inertia plays an extremely important role throughout the wingbeat cycle (Heers et al., 2018). The relative effects of inertia are particularly substantial when aerodynamic force production is low, as during WAIR. This is especially true for immature birds [which produce very small amounts of aerodynamic force (Tobalske & Dial, 2007; Heers et al., 2011)] and presumably for extinct avian predecessors with incipient aerodynamic capacity.

Finally, it is worth noting that the calculations presented by the authors may be influenced by experimental or analytic limitations. For example, it is difficult to determine particular phases of upstroke and downstroke based solely on the figure from which the authors extracted the data used in their analysis. This is because the figure and referenced video frames were likely from different trials and/or individuals. In addition, only one wingbeat was analysed, and at a point when the bird was decelerating. Over many wingbeats, a different pattern might emerge given that the wings and legs oscillate at different frequencies, as the authors have noted. It is also possible that some forces do not act through the centre of mass and therefore produce torques; e.g. a downward-directed wing force helps to pitch the body forward (since the shoulder joints are likely anterior to the centre of mass) and balance the animal on the incline. Regardless, altering the stroke plane angle slightly helps redirect the animal during flight (Dial et al., 2008). Indeed, many birds alter their stroke planes to achieve different flapping behaviours. That WAIR may differ from flight is therefore not unexpected, and neither supports nor precludes WAIR and similar behaviours as potentially important evolutionary drivers.

#### MUSCULOSKELETAL CONSIDERATIONS

In addition to the aerodynamic considerations outlined above, the authors also suggest that WAIR is inconsistent with the evolution of the supracoracoideus muscle, from a 'depressor-protractor' to a 'protractor' (Ostrom, 1976; Novas *et al.*, 2021) and then to an 'elevator and supinator' as well. They state that:

- 1. 'The humeral protractor muscles are not required at all during WAIR'
- 2. "The morphologically necessary protractor state of the supracoracoideus muscle in a "semi-flying" ancestor, which is logically explained by the classical hypothesis of a gliding ancestor, does not fit the WAIR hypothesis of flapping flight origin'
- 3. 'Nowhere in the downstroke of WAIR, [are] the anterior (clavicular) fibres of the pectoralis muscle are required. Therefore, the avian-specific spreading of the pectoralis muscle origin onto the clavicle cannot be explained through the WAIR hypothesis'.

However, all of these statements are inconsistent with previously-published studies. Based on X-ray videos, which provide the most accurate method for quantifying joint movements, protraction is significant (~ 50°) during WAIR (Baier et al., 2013; Heers et al., 2016). Musculoskeletal modelling simulations of WAIR, which align with previously reported data on muscle activity, indicate that both the pectoralis and the supracoracoideus contribute to this protraction (Heers et al., 2018). With respect to the spreading of the pectoralis muscle origin onto the clavicle, it should be noted that changes in bone orientation may have also contributed to this expansion, in addition to locomotor demands. For example, as the sternum expanded and the coracoid became more angled (i.e. producing a smaller scapulocoracoid angle), the furcula may

have expanded ventrally to help maintain the cranial origins of the pectoralis (Heers *et al.*, 2021). Thus, it is not necessary to assume that the spreading of the pectoralis muscle was to enhance protraction—it also could have been a mechanism for evolving a larger and more powerful muscle.

Based on the aerodynamic and musculoskeletal considerations discussed above, the authors conclude that 'WAIR is a very specialized mode of locomotion that is employed by a few specialized birds as an adaptation to a very specific environment and involving highly developed flying features of the locomotor apparatus. Our observations yield a very different conclusion here as well. WAIR and behaviours like WAIR are used by many avian species in many environments (https://youtu.be/VFUNhTdcNdk). Given that WAIR, and similar behaviours like steaming (i.e. using the feet like paddles and the wings like oars to swim) (Dial & Carrier, 2012) or wing-assisted jumping (Heers & Dial, 2015), are extremely common and employed by a diverse array of juvenile birds with very underdeveloped anatomical features compared to adults, it appears to be one of the *least* specialized flapping behaviours observed among extant birds. Indeed, behaviours like WAIR require less aerodynamic force production (Tobalske & Dial, 2007; Heers et al., 2011) and less muscle power (Jackson et al., 2011; Heers et al., 2018), and can be performed by animals that have small wings with less aerodynamically effective feathers (Heers et al., 2011; Dial et al., 2012), smaller muscles (Heers & Dial, 2015), and less robust and specialized skeletons (Heers & Dial, 2012; Heers et al., 2016).

#### LOOKING AHEAD

We heartily agree with Kuznetsov and Panyutina that evolutionary hypotheses should be tested and discussed. However, we believe that some conversations are more productive when they occur directly between scientists as a collaboration, rather than indirectly between articles. We would also like to reiterate that our fields can benefit from:

- 1. *Testing all hypotheses and behaviours similarly*—to keep conclusions in perspective and help validate methods of analysis. For example, some models describing the evolution of avian flight can be applied only to a subset of extant birds or behaviours, and limitations like these should be acknowledged.
- 2. An emphasis on acquiring new data rather than reanalysing old data—but when appropriate, communicating with the authors whose data is being reanalysed.
- 3. Collaborating more across disciplines. As many authors have pointed out, work with extant

organisms is necessary to validate methods of analysing extinct ones (Hutchinson & Allen, 2009; Hutchinson, 2011; Bishop et al., 2021). However, eliciting behaviours and maximal performance in living animals is challenging—should one conclude that a pigeon cannot fly if it does not fly in front of a video camera? Instances like this have occurred. As biologists and palaeobiologists, we probably all agree that extinct animals can provide rich insight into extant ones, and vice versa, and our fields should reflect this belief-by encouraging new ideas, promoting rigorous and interdisciplinary collaborations, and facilitating constructive discussions in welcoming environments.

# ACKNOWLEDGEMENTS

We thank Drs Kutnetzov and Panyutina for raising the issue of WAIR and for their helpful comments on the manuscript, alongside those of two other anonymous referees.

# DATA AVAILABILITY

This article has not generated any new data.

# REFERENCES

- Baier DB, Gatesy SM, Dial KP. 2013. Three-dimensional, high-resolution skeletal kinematics of the avian wing and shoulder during ascending flapping flight and uphill flaprunning. *PLoS One* 8: e63982.
- **Bishop PJ, Cuff AR, Hutchinson JR. 2021.** How to build a dinosaur: musculoskeletal modeling and simulation of locomotor biomechanics in extinct animals. *Paleobiology* **47**: 1–38.
- **Brown RHJ. 1953.** The flight of birds: II. Wing function in relation to flight speed. *Journal of Experimental Biology* **30**: 90–103.
- Bundle MW, Dial KP. 2003. Mechanics of wing-assisted incline running (WAIR). *The Journal of Experimental Biology* 206: 4553–4564.
- Chin DD, Lentink D. 2017. How birds direct impulse to minimize the energetic cost of foraging flight. *Science Advances* 3: e1603041.
- Crandell KE, Tobalske BW. 2011. Aerodynamics of tipreversal upstroke in a revolving pigeon wing. *The Journal of Experimental Biology* 214: 1867–1873.
- Crandell KE, Tobalske BW. 2015. Kinematics and aerodynamics of avian upstrokes during slow flight. *Journal* of Experimental Biology 218: 2518–2527.
- Dial, KP. 2003. Wing-assisted incline running and the evolution of flight. *Science* 299: 402–404.
- Dial TR, Carrier DR. 2012. Precocial hindlimbs and altricial forelimbs: partitioning ontogenetic strategies in mallard

ducks (Anas platyrhynchos). The Journal of Experimental Biology **215:** 3703–3710.

- Dial TR, Heers AM, Tobalske BW. 2012. Ontogeny of aerodynamics in mallards: comparative performance and developmental implications. *The Journal of Experimental Biology* 215: 3693–3702.
- Dial KP, Jackson BE, Segre P. 2008. A fundamental avian wing-stroke provides a new perspective on the evolution of flight. *Nature* **451**: 985–989.
- Dial KP, Randall RJ, Dial TR. 2006. What use is half a wing in the ecology and evolution of birds? *BioScience* 56: 437–445.
- Hedrick TL, Usherwood JR, Biewener AA. 2004. Wing inertia and whole-body acceleration: an analysis of instantaneous aerodynamic force production in cockatiels (*Nymphicus hollandicus*) flying across a range of speeds. Journal of Experimental Biology 207: 1689–1702.
- Heers AM, Baier DB, Jackson BE, Dial KP. 2016. Flapping before flight: high resolution, three-dimensional skeletal kinematics of wings and legs during avian development. *PLoS One* 11: e0153446.
- Heers AM, Dial KP. 2015. Wings versus legs in the avian bauplan: development and evolution of alternative locomotor strategies. *Evolution* **69:** 305–320.
- Heers AM, Dial KP. 2012. From extant to extinct: locomotor ontogeny and the evolution of avian flight. *Trends in Ecology* & *Evolution* 27: 296–305.
- Heers AM, Rankin JW, Hutchinson JR. 2018. Building a bird: musculoskeletal modeling and simulation of wingassisted incline running during avian ontogeny. *Frontiers in Bioengineering and Biotechnology* **6**: 140.
- Heers AM, Tobalske BW, Dial KP. 2011. Ontogeny of lift and drag production in ground birds. *Journal of Experimental Biology* 214: 717–725.
- Heers AM, Varghese SL, Hatier LK, Cabrera JJ. 2021. Multiple functional solutions during flightless to flightcapable transitions. *Frontiers in Ecology and Evolution* 8: 438.
- Hutchinson JR. 2011. On the inference of function from structure using biomechanical modelling and simulation of extinct organisms. *Biology Letters* 8: 115–118.
- Hutchinson JR, Allen V. 2009. The evolutionary continuum of limb function from early theropods to birds. *Naturwissenschaften* 96: 423-448.
- Jackson BE, Tobalske BW, Dial KP. 2011. The broad range of contractile behaviour of the avian pectoralis: functional and evolutionary implications. *The Journal of Experimental Biology* 214: 2354–2361.
- **Kuznetsov AN, Panyutina AA. 2022.** Where was WAIR in avian flight evolution? *Biological Journal of the Linnean Society*, blac019, doi:10.1093/biolinnean/blac019.
- Lorenz K. 1933. Beobachtetes ü ber das Fliegen der Vögel und ü ber die Beziehungen der Flügel- und Steuerform zur Art des Fluges. *Journal of Ornithology* 81: 107–236.
- Marey EJ. 1890. Physiologie du mouvement : Le vol des oiseaux. In : G. Masson ed. Librairie de l'Académie de Médecine, Paris.
- Novas FE, Motta MJ, Agnolín FL, Rozadilla S, Lo Coco GE, Brissón Egli F. 2021. Comments on the morphology of basal paravian shoulder girdle: new data based on unenlagiid

theropods and paleognath birds. Frontiers in Earth Science 9: 386.

- **Ostrom JH. 1976.** Some hypothetical anatomical stages in the evolution of avian flight. *Smithsonian Contributions to Paleobiology* **27:** 1–21.
- Ros IG, Bassman LC, Badger MA, Pierson AN, Biewener AA. 2011. Pigeons steer like helicopters and generate down- and upstroke lift during low speed turns. *Proceedings of the National Academy of Sciences* 108: 19990–19995.
- **Tobalske BW. 2000.** Biomechanics and physiology of gait selection in flying birds. *Physiological and Biochemical Zoology* **73:** 736–750.
- **Tobalske BW, Biewener AA. 2008.** Contractile properties of the pigeon supracoracoideus during different modes of flight. *Journal of Experimental Biology* **211:** 170–179.
- **Tobalske BW, Dial KP. 2000.** Effects of body size on takeoff flight performance in the Phasianidae (Aves). *Journal of Experimental Biology* **203:** 3319–3332.

- Tobalske BW, Dial KP. 2007. Aerodynamics of wing-assisted incline running in birds. *Journal of Experimental Biology* 210: 1742–1751.
- Tobalske BW, Hedrick TL, Biewener AA. 2003. Wing kinematics of avian flight across speeds. *Journal of Avian Biology* 34: 177–184.
- Vazquez RJ. 1992. Functional osteology of the avian wrist and the evolution of flapping flight. *Journal of Morphology* 211: 259–268.
- Vazquez RJ. 1994. The automating skeletal and muscular mechanisms of the avian wing (Aves). Zoomorphology 114: 59–71.
- Warrick DR, Tobalske BW, Powers DR. 2005. Aerodynamics of the hovering hummingbird. *Nature* 435: 1094–1097.
- Warrick DR, Tobalske BW, Powers DR. 2009. Lift production in the hovering hummingbird. *Proceedings of the Royal Society B: Biological Sciences* 276: 3747–3752.